


Lesson Summary

- A rotation of 180 degrees around O is the rigid motion so that if P is any point in the plane, P , O , and $\text{Rotation}(P)$ are *collinear* (i.e., lie on the same line).
- Given a 180-degree rotation, R_0 around the origin O of a coordinate system, R_0 , and a point P with coordinates (a, b) , it is generally said that $R_0(P)$ is the point with coordinates $(-a, -b)$.

THEOREM: Let O be a point not lying on a given line L . Then, the 180-degree rotation around O maps L to a line parallel to L .

Problem Set

Use the following diagram for Problems 1–5. Use your transparency as needed.



1. Looking only at segment BC , is it possible that a 180° rotation would map segment BC onto segment $B'C'$? Why or why not?
2. Looking only at segment AB , is it possible that a 180° rotation would map segment AB onto segment $A'B'$? Why or why not?

- Looking only at segment AC , is it possible that a 180° rotation would map segment AC onto segment $A'C'$? Why or why not?
- Connect point B to point B' , point C to point C' , and point A to point A' . What do you notice? What do you think that point is?
- Would a rotation map triangle ABC onto triangle $A'B'C'$? If so, define the rotation (i.e., degree and center). If not, explain why not.
- The picture below shows right triangles ABC and $A'B'C'$, where the right angles are at B and B' . Given that $AB = A'B' = 1$, and $BC = B'C' = 2$, and that \overline{AB} is not parallel to $\overline{A'B'}$, is there a 180° rotation that would map $\triangle ABC$ onto $\triangle A'B'C'$? Explain.

